



A course on the Ising model





Thursdays 9h45-11h45, Room 15-25.101 (Campus Pierre et Marie Curie)

Start date: 30 January 2025


plammers@lpsm.paris

pietlammers.com#teaching

Pierre Curie observed that a magnet's strength depends on the temperature




Pierre Curie observed that a magnet's strength depends on the temperature



- What mechanism changes the strength?
- What happens at the **critical point**?
- What happens around the **critical point**?

Pierre Curie observed that a magnet's strength depends on the temperature




- What mechanism changes the strength?
- What happens at the **critical point**?
- What happens around the **critical point**?

Mathematically, $M(T) = 0$ for $T > T_c$,
 $M(T) > 0$ for $T < T_c$;

Loss of **regularity** of M at $T=T_c$.

Pierre Curie observed that a magnet's strength depends on the temperature



- What mechanism changes the strength?
- What happens at the **critical point**?
- What happens around the **critical point**?



Mathematically, $M(T) = 0$ for $T > T_c$,
 $M(T) > 0$ for $T < T_c$;

Loss of **regularity** of M at $T=T_c$.



Phase transition

Lenz proposed a mathematical model (~1920)
to his PhD student to “simulate” magnets:
the **Ising model**

Lenz proposed a mathematical model (~ 1920)
to his PhD student to “simulate” magnets:
the **Ising model**





Lenz proposed a mathematical model (~1920) to his PhD student to “simulate” magnets: the **Ising model**



Ising solved the model in **one dimension** (one hundred years ago, in 1925). With the knowledge of today, this is straightforward because we have the theory of **Markov chains**.

Lenz proposed a mathematical model (~1920) to his PhD student to “simulate” magnets: the **Ising model**



Ising: “The model has no phase transition!”

Ising solved the model in **one dimension** (one hundred years ago, in 1925). With the knowledge of today, this is straightforward because we have the theory of **Markov chains**.

Lenz was right in 1D, but higher dimensions is much more complicated!

Complications arise because:

- 1D is straightforward because of Markov chains, but
- In higher dimensions, the geometry plays a huge role!


Example:

- Random walk is recurrent in 2D, but transient in 3D and higher D,
- Paths of random walks intersect in 3D, but not in 4D and higher D,


The Ising model:

- Behaves very differently in dimensions 1, 2, 3, 4, and 5+

The Ising model per dimension



- 1D: Trivial, Markov chains
- 2D: Almost everything is known (integrability, transfer matrix)
- 3D: Almost nothing is known
- 4D: Many things are known (random walks behave well)
- 5D: Almost everything is known (random walks behave very well)



Objectives of the course:

- Overview of **statistical mechanics (goals, ideas, methods)**
- Know the **key tools** and how to apply them
- Have an **historical overview** of the developments



Mathematically:

- Existence & uniqueness of the phase transition (1936 Peierls argument)
- Sharpness of the phase transition
- Continuity of the phase transition (2022 Fields medal, Duminil-Copin)

(Pre)requisites:

- Basic understanding of probability theory and graph theory
- The course will be reasonably demanding:
 - Some time to reflect on the material in between lectures will be useful

A course on the Ising model



Thursdays 9h45-11h45, Room 15-25.101 (Campus Pierre et Marie Curie)

Start date: 30 January 2025

`plammers@lpsm.paris`

`pietlammers.com#teaching`

QUESTIONS WELCOME!