

# Conformal invariance of the planar four-state Potts model and its random-cluster representation

Emile Averous\*, Hugo Duminil-Copin†, Tiancheng He‡  
Piet Lammers§ and Ioan Manolescu¶

6 October 2026

## Abstract

For the critical random-cluster model on  $\mathbb{Z}^2$  with cluster-weight  $q = 4$ , we prove the convergence of loops to  $\text{CLE}(4)$  in the scaling limit. Conformal invariance of the four-state Potts model is then deduced via the Edwards–Sokal coupling.




Figure 1: **The critical random-cluster model at  $q = 4$ .** One may view the model as follows: fix a deterministic boundary loop on the medial lattice of  $\mathbb{Z}^2$  (a scaled, rotated version of the square lattice), and sample, uniformly at random, a collection of directed non-intersecting loops inside the domain, occupying all edges of the medial lattice. In this simulation, we highlighted the eight largest loops (in terms of enclosed area). Our result asserts that, in the scaling limit, the loops' law converges to  $\text{CLE}(4)$ .

---

\*Université de Genève, [emile.averous@unige.ch](mailto:emile.averous@unige.ch)

†Université de Genève, [hugo.duminil@unige.ch](mailto:hugo.duminil@unige.ch), Institut des Hautes Études Scientifiques, [duminil@ihes.fr](mailto:duminil@ihes.fr)

‡Université de Genève, [tiancheng.he@unige.ch](mailto:tiancheng.he@unige.ch)

§CNRS, Sorbonne Université, LPSM, [piet.lammers@cnrs.fr](mailto:piet.lammers@cnrs.fr)

¶Université de Fribourg, [ioan.manolescu@unifr.ch](mailto:ioan.manolescu@unifr.ch)

# 1 Introduction

## 1.1 Motivation

Conformal Field Theory (CFT) has been a cornerstone of theoretical physics in the twentieth century [6, 7]. The introduction of the Schramm–Loewner Evolution (SLE) [45, 33] and Conformal Loop Ensembles (CLE) [49, 50] (see also [36, 37]), together with major advances in the mathematical foundations of CFT [17, 44, 28], has profoundly reshaped our understanding of conformally invariant objects and two-dimensional CFTs.

Despite this progress, rigorous proofs of conformal invariance for lattice models remain remarkably scarce. Among the principal examples are: Bernoulli site percolation on the triangular lattice [51, 32, 11, 10] (results range from Cardy’s formula to convergence of interfaces and loop ensembles toward  $\text{SLE}_6$  and  $\text{CLE}_6$ ); the Ising and FK-Ising models [52, 13, 29, 12, 14, 8] (major breakthroughs include the analysis of fermionic observables, energy and spin fields, and convergence toward SLE and CLE with parameters 3 and  $16/3$ ); uniform spanning trees [34]; dimer models [30, 31]; and level lines of the discrete Gaussian free field [46].

More recently, progress towards a more systematic analysis of scaling limits of lattice models was achieved with the proof of conformal invariance of the six-vertex model for a continuum of parameters; see [23] for the full-plane result and [2] for the corresponding result in simply connected domains. In the present paper, we build on [2] to establish conformal invariance of the random-cluster model with cluster weight  $q = 4$  on the square lattice, as well as of the associated four-state Potts model. This represents a special case of the Ashkin-Teller model, which we plan to treat in the subsequent paper [3], generalizing the arguments in [22, 21].

## 1.2 Definition of the random-cluster model

Consider a finite subgraph  $G = (V, E)$  of  $\mathbb{Z}^2$ . A percolation configuration  $\omega$  on  $G$  is an element of  $\{0, 1\}^E$ . We say that an edge  $e$  is *open* if  $\omega(e) = 1$  and that it is *closed* if  $\omega(e) = 0$ . We can see a configuration  $\omega$  as a subgraph of  $G$  with the vertex-set  $V$  and the edge-set  $\{e \in E : \omega(e) = 1\}$ . A *cluster* is a connected component of  $\omega$  (possibly including isolated vertices).

**Definition 1.1.** *The random cluster measure on  $G$  with edge-weight  $p \in (0, 1)$ , cluster-weight  $q > 0$  and free boundary conditions is given by*

$$\phi_{G,p,q}^0[\omega] := \frac{1}{Z^0(G, p, q)} \left( \frac{p}{1-p} \right)^{|\omega|} q^{k(\omega)},$$

where  $|\omega| := \sum_{e \in E} \omega(e)$ ,  $k(\omega)$  is the number of connected components of the graph  $\omega$  that intersect  $V$ , and  $Z^0(G, p, q)$  is the partition function making  $\phi_{G,p,q}^0$  a probability measure.

Consider the dual graph  $(\mathbb{Z}^2)^* := (\frac{1}{2}, \frac{1}{2}) + \mathbb{Z}^2$ . Note that every edge  $e$  of  $\mathbb{Z}^2$  has a dual-edge  $e^*$  of  $(\mathbb{Z}^2)^*$  crossing it in its middle. For a percolation configuration  $\omega$  on  $\mathbb{Z}^2$ , let  $\omega^*$  be the configuration on  $(\mathbb{Z}^2)^*$  defined by the formula  $\omega^*(e^*) = 1 - \omega(e)$  for every edge  $e$ . In words, a dual-edge is open if the corresponding edge of  $\mathbb{Z}^2$  is closed, and vice-versa.

Consider now the medial graph  $(\mathbb{Z}^2)^\diamond$  of the square lattice obtained by putting a medial vertex at the middle of each edge of  $\mathbb{Z}^2$ , and medial edges between nearest medial




Figure 2: **Coupling of several models.** The **random-cluster model** is a percolation model on the square lattice  $\mathbb{Z}^2$ . This simulation has free boundary conditions. The **Potts model** is obtained by colouring its clusters uniformly at random with four colours. The **loops** live on the medial lattice and separate the primal clusters from the dual clusters (the dual clusters are not drawn). They are given an *orientation* by flipping a fair coin for each loop. Notice that the random-cluster model and the *unoriented* loops encode precisely the same information.

vertices. The graph is a rescaled (by a factor  $1/\sqrt{2}$ ) and rotated (by an angle  $\pi/4$ ) version of the square lattice. For a configuration  $\omega$  on  $\mathbb{Z}^2$ , there is a unique way to introduce a loop configuration  $\bar{\omega}$  on  $(\mathbb{Z}^2)^\diamond$  in such a way that loops are drawn in the medial lattice, they turn at each medial vertex so that they do not cross any open edge of  $\omega$  or  $\omega^*$ , and all medial edges are part of exactly one loop; see Figure 2.

In what follows, we call *discrete domain* any subgraph  $G$  of  $\mathbb{Z}^2$  obtained by taking all the vertices (of  $\mathbb{Z}^2$ ) within a loop  $\ell$  on the medial lattice  $(\mathbb{Z}^2)^\diamond$  turning at every vertex, and all edges (of  $\mathbb{Z}^2$ ) between these vertices not crossing the loop  $\ell$ . A discrete domain can be seen as a union of simply connected open sets of  $\mathbb{R}^2$  by considering the points enclosed by the loop<sup>1</sup>. The boundary  $\partial G$  of  $G$  is the set of vertices in  $V$  that correspond to (medial) faces of  $(\mathbb{Z}^2)^\diamond$  bordering the loop  $\ell$ .

We consider a percolation configuration  $\omega$  on a finite graph  $G = (V, E)$  with free boundary conditions as a configuration on the whole lattice  $\mathbb{Z}^2$  by declaring all edges which are not in  $E$  to be closed. As a consequence, we may define  $\bar{\omega}$  by considering the full-plane configuration and then removing all the small loops surrounding vertices which are not in  $V$ . With this definition, the configuration  $\bar{\omega}$  is composed entirely of loops.

For more details on the random-cluster model, see e.g. [27, 19].

*In this article, we focus on the cluster weight  $q = 4$ . In such a case, it has been proved*

<sup>1</sup>The loops are considered as making small roundish turns at every medial vertex (cf. Figure 1).

that the model undergoes a continuous phase transition at the self-dual point  $p_c = 2/3$  [26]. In all that follows, we set  $p := p_c = 2/3$ .

### 1.3 Conformal invariance of the random-cluster model at $q = 4$

The object of interest in the continuum will be the nested Conformal Loop Ensemble (CLE) with parameter  $\kappa$  equal to 4. We refer to [49, 50] for its definition. Here, we simply recall that the CLE(4) in a simply connected domain  $\mathcal{D}$  is a random collection of nested, non intersecting and non self-crossing continuous loops in  $\mathcal{D}$  which is conformally invariant, in the sense that CLE(4) in  $\Psi(\mathcal{D})$ , where  $\Psi$  is a conformal map, is the image by  $\Psi$  of CLE(4) in  $\mathcal{D}$ .

We say that a family of discrete domains  $\mathcal{D}_\delta$  on  $\delta\mathbb{Z}^2$  converges to  $\mathcal{D}$  in the *Carathéodory sense* if there exist conformal bijections  $f_\delta : \mathcal{D}_\delta \rightarrow \mathbb{D}$  and  $f : \mathcal{D} \rightarrow \mathbb{D}$  such that every compact subset of  $\mathcal{D}$  is contained in  $\mathcal{D}_\delta$  for all sufficiently small  $\delta$ , and  $f_\delta$  converges to  $f$  uniformly on every compact subset of  $\mathcal{D}$ .

**Theorem 1.2** (CLE(4) convergence of macroscopic loops (Figure 1)). *Consider a simply connected domain  $\mathcal{D}$  and let  $(\mathcal{D}_\delta)$  be a family of discrete domains on  $\delta\mathbb{Z}^2$  converging to  $\mathcal{D}$  in the Carathéodory sense. Let  $\bar{\omega}_\delta$  be the loop configuration obtained from the random-cluster model on  $\mathcal{D}_\delta$  with cluster-weight  $q = 4$  and edge-weight  $p = 2/3$ , with free boundary conditions. Then,  $\bar{\omega}_\delta$  converges weakly (for the topology induced by the Camia-Newman distance<sup>2</sup>) to the CLE(4) in  $\mathcal{D}$ .*

Using the universality result of [24], the theorem extends to every rectangular isoradial graph. We chose to state it in the case of the square lattice for simplicity. Previous results concerning the random-cluster model with cluster-weight  $q = 4$  include the derivation of the critical exponents [15], as well as certain correlations in full plane [16]. Both use the GFF convergence of [23].

The following corollary, based on our main result combined with mixing properties of the model, states that the multi-point connectivity correlations are conformally invariant. The exact expression of these correlations can be obtained in terms of CLE(4) correlations, but we do not include it here. Below, for a point  $x \in \mathbb{R}^2$ , let  $[x]_\delta$  be the vertex of  $\delta\mathbb{Z}^2$  closest to  $x$  (in case of a tie we choose one of the closest vertices according to an arbitrary rule).

---

<sup>2</sup>For a simply connected domain  $\Omega$ , let  $\mathfrak{C} = \mathfrak{C}(\Omega)$  be the collection of locally finite families  $\mathcal{F}$  of non-self-crossing loops in  $\Omega$  that do not intersect each other. Define the metric on  $\mathfrak{C}$ ,

$$d_{\text{CN}}(\mathcal{F}, \mathcal{F}') \leq \varepsilon \iff \left( \begin{array}{l} \forall \gamma \in \mathcal{F} \text{ with } \gamma \subset B(0, 1/\varepsilon), \exists \gamma' \in \mathcal{F}', d(\gamma, \gamma') \leq \varepsilon \\ \text{and similarly when exchanging } \mathcal{F}' \text{ and } \mathcal{F} \end{array} \right),$$

where, for two loops  $\gamma_1$  and  $\gamma_2$ , we set

$$d(\gamma_1, \gamma_2) := \inf_{t \in \mathbb{S}^1} \sup_{t \in \mathbb{S}^1} |\gamma_1(t) - \gamma_2(t)|,$$

with the infimum running over all continuous one-to-one parametrizations of the loops  $\gamma_1$  and  $\gamma_2$  by  $\mathbb{S}^1$ .

The distance  $d_{\text{CN}}$  induces a distance on the space of probability measures on  $\mathfrak{C}$ , which we also denote by  $d_{\text{CN}}$ : for  $\phi$  and  $\phi'$  two measures on  $\mathfrak{C}$ , let

$$d_{\text{CN}}(\phi, \phi') := \inf \{ \exists \text{ coupling } \mathbb{P} \text{ of } \phi \text{ and } \phi' \text{ with } \mathbb{P}[d_{\text{CN}}(\bar{\omega}, \bar{\omega}') > \varepsilon] < \varepsilon \}.$$

**Corollary 1.3** (Conformal invariance of point-to-point correlations). *There exists  $\kappa(\delta)$  such that  $\lim_{\delta \rightarrow 0} \frac{\log(\kappa(\delta))}{\log(1/\delta)} = 1/8$  and the following holds. Consider a simply connected domain  $\mathcal{D}$  and  $n \geq 1$ . For any partition  $P = P_1 \sqcup \dots \sqcup P_s$  of  $\{1, \dots, n\}$ , there exists a map  $\Psi_{\mathcal{D}}(P; \cdot)$  such that for every  $x_1, \dots, x_n \in \mathcal{D}$ ,*

$$\lim_{\delta \rightarrow 0} \kappa(\delta)^n \phi_{\mathcal{D}, 2/3, 4}^0[\mathcal{E}_{\delta}(P; x_1, \dots, x_n)] = \Psi_{\mathcal{D}}(P; x_1, \dots, x_n),$$

where  $\mathcal{E}_{\delta}(P; x_1, \dots, x_n)$  denotes the event that  $[x_i]_{\delta}$  and  $[x_j]_{\delta}$  are connected in  $\omega_{\delta}$  if and only if  $i$  and  $j$  are in the same  $P_t$  for some  $1 \leq t \leq s$ .

Furthermore, the family of functions  $(\Psi_{\mathcal{D}}(P; \cdot))_{\mathcal{D}}$  is conformally covariant in the following sense: for every conformal map  $f : \mathcal{D} \rightarrow \mathbb{C}$  and every  $x_1, \dots, x_n \in \mathcal{D}$ ,

$$\Psi_{\mathcal{D}}(P; x_1, \dots, x_n) = |f'(x_1)|^{1/8} \dots |f'(x_n)|^{1/8} \Psi_{f(\mathcal{D})}(P; f(x_1), \dots, f(x_n)).$$

## 1.4 Conformal invariance of the Potts model

For a finite subgraph  $G = (V, E)$  of  $\mathbb{Z}^2$ , attribute a *spin* variable  $\sigma_x$  belonging to  $\{1, \dots, q\}$  to each vertex  $x \in V$ . A *spin configuration*  $\sigma = (\sigma_x : x \in V) \in \{1, \dots, q\}^V$  is given by the collection of all the spins. Introduce the Hamiltonian of  $\sigma$  defined by

$$H_G(\sigma) := - \sum_{xy \in E} \mathbb{1}[\sigma_x = \sigma_y].$$

The *Gibbs measure* on  $G$  at inverse temperature  $\beta \geq 0$  with free boundary conditions is defined by the formula

$$\mu_{G, \beta, q}[\sigma] := \frac{1}{Z(G, \beta, q)} \exp[-\beta H_G(\sigma)],$$

for every  $\sigma \in \{1, \dots, q\}^V$ , and  $Z(G, \beta, q)$  is the partition function making the previous measure a probability measure.

This model was introduced as a generalization of the Ising model to more than two possible spins by Potts in 1952 [42] following a suggestion of his adviser Domb. While the model received little attention early on, it became an object of great interest in the last forty years. Since then, mathematicians and physicists have been studying it intensively, and a lot is known on its rich behavior.

The Potts model with  $q$  states is related to the random-cluster model with cluster-weight  $q$  via the Edwards–Sokal coupling ; see e.g. [27, 19]. The Edwards–Sokal coupling has many applications. Here, we mention the simplest statement, and leave more evolved statements to the reader. Below, we focus on the case  $q = 4$ . It is easy to deduce from the critical point of the random-cluster model that the 4-state Potts model undergoes a continuous phase transition at  $\beta_c = \log(3)$ . Recall that  $\kappa(\delta)$  is given by Corollary 1.3 and satisfies  $\lim_{\delta \rightarrow 0} \frac{\log(\kappa(\delta))}{\log(1/\delta)} = 1/8$ .

**Corollary 1.4.** *Consider a simply connected domain  $\mathcal{D}$ . Fix a tuple of spins  $\tau \in \{1, \dots, q\}^n$ . There exists  $\tilde{\Psi}_{\mathcal{D}}(\tau; \cdot)$  such that for every  $x_1, \dots, x_n \in \mathcal{D}$ ,*

$$\lim_{\delta \rightarrow 0} \kappa(\delta)^n \mu_{\mathcal{D}, \beta_c, 4} \left[ \prod_{i=1}^n (\mathbb{1}[\sigma_{[x_i]_{\delta}} = \tau_i] - \frac{1}{4}) \right] = \tilde{\Psi}_{\mathcal{D}}(\tau; x_1, \dots, x_n).$$

Furthermore, the family of functions  $(\tilde{\Psi}_{\mathcal{D}}(\tau; \cdot))_{\mathcal{D}}$  is conformally covariant in the following sense: for every conformal map  $f : \mathcal{D} \rightarrow \mathbb{C}$  and every  $x_1, \dots, x_n \in \mathcal{D}$ ,

$$\tilde{\Psi}_{\mathcal{D}}(\tau; x_1, \dots, x_n) = |f'(x_1)|^{1/8} \dots |f'(x_n)|^{1/8} \tilde{\Psi}_{f(\mathcal{D})}(\tau; f(x_1), \dots, f(x_n)).$$




Figure 3: The **six-vertex configuration** is obtained from the *oriented* loops by “forgetting” the way that the oriented edges are wired up into loops; see Figure 2 for the same simulation *with* the orientations.

## 2 Proof of Theorem 1.2

The proof relies on a coupling between the random-cluster model and the height function of the six-vertex model. We start by defining the six-vertex model. Then, we recall the convergence result obtained in [2]. Finally, we prove our result.

### 2.1 The six-vertex model

The six-vertex model is a probability measure on orientations of the edges of a graph [4, 35]. A six-vertex configuration  $\omega^\times$  on a subgraph of the medial lattice  $(\mathbb{Z}^2)^\diamond$  assigns an orientation to each edge, subject to the *ice rule*: at every vertex of degree four, the number of incoming edges equals the number of outgoing edges. In particular, there are precisely six local arrow configurations compatible with this constraint, as depicted in Figure 4. The six local configurations are grouped into three pairs, to which we assign positive weights **a**, **b**, and **c**, referred to below as *Boltzmann weights*.

The six-vertex model is often considered with periodic boundary conditions. In this paper, however, we work exclusively in finite domains with fixed boundary conditions. For a discrete domain  $G$ , let  $G^\diamond$  be the collection  $F$  of faces of  $(\delta\mathbb{Z}^2)^\diamond$  which are enclosed by the medial loop defining the boundary of  $G$ . Note that the degree of vertices in  $G^\diamond$  is two or four.

Every six-vertex configuration  $\omega^\times$  on  $G^\diamond$  naturally gives rise to an integer-valued *height function*  $h$  defined on  $F$ . It is piecewise constant on each face and its increments between neighboring faces are determined by the orientation of the edge separating them: when crossing an edge of  $\omega^\times$ , the height increases by 1 if the crossed arrow points to the right




Figure 4: The six possible local arrow configurations around a square lattice vertex of the six-vertex model. These local configurations get the weights  $\mathbf{a}$ ,  $\mathbf{a}$ ,  $\mathbf{b}$ ,  $\mathbf{b}$ ,  $\mathbf{c}$ ,  $\mathbf{c}$  respectively. The  $\mathbf{c}$ -type vertices are special; they correspond to saddle points of the height function.

and decreases by 1 otherwise.

The ice rule ensures that these local increments are compatible around every vertex and can therefore be integrated to define a height function on  $F$ . The height function is determined only up to a global additive constant, whereas its gradient is uniquely determined by  $\omega^\times$ . Conversely, the arrow configuration can be recovered from the height increments between neighboring faces. Thus, modulo an additive constant, the height function and the six-vertex configuration encode the same information.

Consider a discrete domain  $G$  and recall the definition of  $\partial G$ , i.e. the vertices of  $\mathbb{Z}^2$  (and therefore the associated medial faces) that are adjacent to the medial loop defining the boundary of  $G$ . The six-vertex measure on  $G^\circ$  with *Dirichlet boundary condition 0* is defined by

$$\mathbb{P}_{G,\mathbf{a},\mathbf{b},\mathbf{c}}^0[\omega^\times] := \frac{1}{Z_{G,\mathbf{a},\mathbf{b},\mathbf{c}}^0} \mathbf{a}^{N_1+N_2} \mathbf{b}^{N_3+N_4} \mathbf{c}^{N_5+N_6} \mathbb{1}_{h|_{\partial G}=0}, \quad (1)$$

where  $N_i = N_i(\omega^\times)$  denotes the number of degree-four vertices whose local arrow configuration is of type  $i$ . The normalizing constant  $Z_{G,\mathbf{a},\mathbf{b},\mathbf{c}}^0$  is the corresponding *partition function*, chosen so that (1) defines a probability measure.

*Below, we restrict to the parameters  $\mathbf{a} = \mathbf{b} = 1$  and  $\mathbf{c} = 2$ .*

## 2.2 Scaling limit of the six-vertex height function

Consider a simply connected domain  $\mathcal{D}$ . Let  $G_{\mathcal{D}} : D_2(\mathcal{D}) := \{(u, v) \in \mathcal{D}^2 : u \neq v\} \rightarrow \mathbb{R}$  denote the Dirichlet Green function in  $\mathcal{D}$  defined as follows: for each fixed  $y \in \mathcal{D}$ , the function  $x \mapsto G_{\mathcal{D}}(x, y)$  is harmonic on  $\mathcal{D} \setminus \{y\}$ , vanishes as  $x$  approaches  $\partial\mathcal{D}$ , and satisfies

$$G_{\mathcal{D}}(x, y) = -\frac{1}{2\pi} \log |x - y| + O(1) \quad \text{as } x \rightarrow y.$$

By a *finite-Dirichlet-energy generalized test function* we mean a finite signed measure  $\varphi$  supported in  $\mathcal{D}$  such that

$$\iint_{\mathcal{D} \times \mathcal{D}} G_{\mathcal{D}}(u, v) d\varphi(u) d\varphi(v) < \infty.$$

The Gaussian Free Field (GFF) is a centered Gaussian generalized function  $\Gamma$  with covariance kernel  $G_{\mathcal{D}}$ ; see [9, 48]. More precisely, it is a random distribution, in the sense that it can be tested against a finite-Dirichlet-energy generalized test function  $\varphi$  by setting

$$\langle \Gamma, \varphi \rangle := \int_{\mathcal{D}} \Gamma(u) d\varphi(u).$$

Then, for every finite family  $\varphi = (\varphi_1, \dots, \varphi_n)$  of finite-Dirichlet-energy generalized test functions,

$$(\langle \Gamma, \varphi_i \rangle)_{i=1}^n \sim \mathcal{N}(0, \Sigma(\varphi)),$$

where  $\Sigma(\varphi)$  is the  $n \times n$  matrix defined by

$$\Sigma(\varphi)_{ij} := \iint_{\mathcal{D} \times \mathcal{D}} G_{\mathcal{D}}(u, v) d\varphi_i(u) d\varphi_j(v).$$

The following convergence result for the six-vertex height function was obtained in [2] and is the basis of our proof.

**Theorem 2.1** (GFF convergence of the six-vertex height function [2]). *Consider the six-vertex model with weights  $\mathbf{a} = \mathbf{b} = 1$  and  $\mathbf{c} = 2$ . Let  $\mathcal{D}$  be a simply connected domain. For a family  $(\mathcal{D}_\delta)$  of discrete domains converging to  $\mathcal{D}$  in the Carathéodory sense, the law of the height function  $h_\delta \sim \mathbb{P}_{\mathcal{D}_\delta^\circ, 1, 1, 2}^0$  of the six-vertex model associated with  $\omega \sim \phi_{\mathcal{D}_\delta, 2/3, 4}^0$ , converges to the GFF  $\sigma \cdot \Gamma$  with zero boundary conditions on  $\mathcal{D}$ , and variance  $\sigma^2 = \frac{2}{\pi}$ .*

## 2.3 The Baxter–Kelland–Wu correspondence

There is a formal correspondence, called the Baxter–Kelland–Wu (BKW) correspondence, between the loop configurations and the arrow configurations. We refer to [5] or the more recent expositions [18, 20, 1] for details. In the case of  $\mathbf{a} = \mathbf{b} = 1$  and  $\mathbf{c} = 2$ , it is a probabilistic coupling between the six-vertex model and the random-cluster model, that we now state.

Consider a discrete domain  $G$ . Consider the uniform measure  $\vec{\mathbb{P}}_G$  on oriented loop configurations  $\vec{\omega}$  on  $G^\circ$ , i.e. loop configurations  $\vec{\omega}$  for which each loop receives an orientation, clockwise or counterclockwise. Then, the BKW coupling states the following two observations.

- If  $\bar{\pi}$  denotes the projection from oriented loop configurations to loop configurations consisting in forgetting orientations of loops, then

$$\phi_{G, 2/3, 4}^0 = \bar{\pi}_* \vec{\mathbb{P}}_G.$$

(See Figure 2.)

- If  $\vec{\pi}$  denotes the projection from oriented loop configurations to arrow configurations consisting in forgetting the loops and simply keeping the orientations of edges induced by the orientations of the loops<sup>3</sup>, and setting the height function to be zero on  $\partial G$ , then

$$\mathbb{P}_{G, 1, 1, 2}^0 = \vec{\pi}_* \vec{\mathbb{P}}_G.$$

(See Figure 3.)

## 2.4 Proof of Theorem 1.2

The next lemma states a natural decomposition coming from the coupling between the height function and the loop configuration.

---

<sup>3</sup>Note that when considering the BKW coupling starting from a configuration  $\omega \sim \phi_{G, 2/3, 4}^0$ , the corresponding boundary conditions of the six-vertex height function are constant in the following sense: the height function restricted to medial faces corresponding to vertices of  $\partial G$ , is constant.




Figure 5: **A discrete version of the two-valued local set of Lemma 2.2:** The decomposition is obtained by exploring the outermost loops in the domain. The orientations of the loops are independent fair coin flips. By the Markov property, the law inside each loop is given by an independent copy of the height function with zero boundary conditions, plus the coin flip coming from the orientation of that loop.

In the proof of Theorem 1.2, we argue that this discrete decomposition commutes with the scaling limit and yields the appropriate two-valued local set of the GFF  $\sigma \cdot \Gamma$ , which enables us to identify the scaling limit of the outermost loops as CLE(4).

**Lemma 2.2** (Discrete two-valued local set (cf. Figure 5)). *Consider a discrete domain  $G$ , and let  $h_G$  be the six-vertex height function in  $G$  with zero boundary conditions and variance  $\sigma^2 = \frac{2}{\pi}$ , let  $\mathcal{L} = \mathcal{L}(\bar{\omega})$  be the collection of exterior-most loops in  $\bar{\omega}$ , and let  $\mathcal{D}(\ell)$  be the discrete domain obtained by considering all the medial faces fully enclosed by  $\ell$ . We have the following identity in law:*

$$h_G = \sum_{\ell \in \mathcal{L}(\bar{\omega})} (\xi_\ell \mathbb{1}_{\mathcal{D}(\ell)} + h_{\mathcal{D}(\ell)}), \quad (2)$$

where the  $\xi_\ell$  are iid  $\pm 1$  Bernoulli variables, and the  $h_{\mathcal{D}(\ell)}$  are independent copies of the six-vertex height function in  $\mathcal{D}(\ell)$  with zero boundary conditions.

*Proof.* The decomposition is immediate by the BKW coupling. The variable  $\xi_\ell$  is equal to  $+1$  if the loop  $\ell$  is oriented counterclockwise, and  $-1$  if it is oriented clockwise.  $\square$

Let us remark that in Lemma 2.2, the jumps of the height function from one side of the loop to another are equal to  $\pm 1$ . In the corresponding GFF limit, the jumps of the GFF  $\sigma \cdot \Gamma$  when traversing a loop of the CLE(4) are  $\pm \sigma \cdot 2\sqrt{\frac{\pi}{8}} = \pm 1$  [46], so this discrete decomposition has an exact continuum counterpart. We are now in a position to prove Theorem 1.2.

*Proof of Theorem 1.2.* It was proved in [47] that there is a unique thin local set with the property that the GFF  $\sigma \cdot \Gamma$  has boundary values  $\pm 1$  on the boundary of each connected

component of the complement of the local set. Moreover, this thin local set is precisely the collection of outermost loops of a CLE(4). Below, we argue that the decomposition obtained in Lemma 2.2 commutes with the limit, and yields precisely this thin local set. This proves that the limit of the law of the outermost discrete loops is precisely CLE(4).

We start by studying the scaling limit of exterior-most loops. RSW theory [26] implies precompactness of the family of loops. Thanks to this precompactness, it suffices to show that any sub-sequential limit  $\mathcal{L}$  of the family of loops  $\mathcal{L}(\bar{\omega}_\delta)$  is equal to the exterior-most loops of a CLE(4). Consider such a sub-sequential limit  $\mathcal{L}$ . Let  $\mathcal{A}$  be the “gasket” of the loops in  $\mathcal{L}$ , meaning the closure of the connected component in  $\mathcal{D} \setminus \bigcup_{\ell \in \mathcal{L}} \ell$  of  $\partial\mathcal{D}$ .

First, note that  $\mathcal{A}$  is thin. Indeed, standard RSW estimates [26] imply that for a ball of size  $\varepsilon > 0$  at distance  $r > \varepsilon$  of  $\partial\mathcal{D}$  to be contained in  $\mathcal{A}$ , there should be no loop in  $\mathcal{L}$  surrounding it. This occurs with probability  $(\varepsilon/r)^c$  for some constant  $c > 0$ . The proof of thinness then follows from the same computation as in the proof of [21, Lemma 6.10].

Second, we claim that  $\mathcal{A}$  is a local set in the sense of [47]. Indeed, by the Skorokhod representation theorem assume that the  $\omega_\delta$  are coupled so that the  $\mathcal{L}(\bar{\omega}_\delta)$  converge almost surely to  $\mathcal{L}$ . Using precompactness for the height function seen as a random distribution, we may extract a sub-sequential limit in which  $h_{\mathcal{D}_\delta}$  and  $(h_{\mathcal{D}(\ell_\delta)} : \ell_\delta \in \mathcal{L}(\bar{\omega}_\delta))$  converge almost surely to  $h_{\mathcal{D}}$  and  $(h_{\mathcal{D}(\ell)} : \ell \in \mathcal{L})$ , in the sense that for all loops  $\ell \in \mathcal{L}$ , if a sequence of loops  $\ell_\delta$  of  $\mathcal{L}(\bar{\omega}_\delta)$  converges to  $\ell$ , then  $h_{\mathcal{D}(\ell_\delta)}$  converges to  $h_{\mathcal{D}(\ell)}$  (in the sense of Theorem 2.1). From (2), we deduce the following identity in law

$$h_{\mathcal{D}} = \sum_{\ell \in \mathcal{L}} (\xi_\ell \mathbb{1}_{\mathcal{D}(\ell)} + h_{\mathcal{D}(\ell)}).$$

As a consequence,  $\mathcal{A}$  is indeed a local set. Note that we use uniform thinness (as  $\delta$  tends to zero) of the gaskets of  $\mathcal{L}(\bar{\omega}_\delta)$  (which is obtained by RSW theory [26] in the same way as for  $\mathcal{L}$ ) to justify that small loops in  $\mathcal{L}(\bar{\omega}_\delta)$  (i.e. loops of size  $o(1)$ ) do not contribute in the limit.

Third, we identify the values of the field on the boundary of  $\mathcal{A}$ . Inside each component of  $\mathcal{A}^c$ , the limit height function is a GFF with boundary conditions  $\xi_\ell \in \{-1, +1\}$ , where  $\ell$  is the loop tracing out the boundary of this component. Note that we do not claim that the boundary conditions in each such component are i.i.d. since one loop  $\ell$  can a priori carve multiple components if it touches itself, but it will suffice for us that the boundary conditions are either  $-1$  or  $+1$ .

By the jump calculation preceding the proof and the characterization above, it follows that  $\mathcal{L}$  is the set of exterior most loops of CLE(4); see [47].

The nested result readily follows by the Markov property and precompactness.  $\square$

### 3 Proof of the corollaries

*Proof of Corollary 1.3.* Consider a simply connected domain  $\mathcal{D}$  and  $n \geq 1$ . For any partition  $P = P_1 \sqcup \dots \sqcup P_s$  of  $\{1, \dots, n\}$ , for every  $x_1, \dots, x_n \in \mathcal{D}$  and  $\varepsilon > 0$ , the CLE(4) convergence implies that

$$\lim_{\delta \rightarrow 0} \phi_{\mathcal{D}_\delta, 2/3, 4}^0[\mathcal{E}_\delta^\varepsilon(P; x_1, \dots, x_n)] = \mathbb{P}_{\text{CLE}(4)}(\mathcal{E}^\varepsilon(P; x_1, \dots, x_n)),$$

where  $\mathcal{E}_\delta^\varepsilon(P; x_1, \dots, x_n)$  denotes the event that the Euclidean balls of size  $\varepsilon$  around  $x_i$  and  $x_j$  are connected in  $\omega_\delta$  (resp. the complement of loops in CLE(4)) if and only if  $i$  and  $j$  are in the same  $P_t$ .

The mixing property [15] implies that for every  $\eta > 0$ , there exists  $\varepsilon > 0$  such that

$$\left| \phi_{\mathcal{D}_\delta, 2/3, 4}^0[\mathcal{E}_\delta(P; x_1, \dots, x_n)] - \phi_{\delta\mathbb{Z}^2, 2/3, 4}^0[0 \leftrightarrow \infty | B_\varepsilon \leftrightarrow \infty]^n \phi_{\mathcal{D}_\delta, 2/3, 4}^0[\mathcal{E}_\delta^\varepsilon(P; x_1, \dots, x_n)] \right| \leq \eta.$$

Above, we used the measure  $\phi_{\delta\mathbb{Z}^2, 2/3, 4}^0[\cdot | B_\varepsilon \leftrightarrow \infty]$  which is the limit (as  $R$  tends to infinity) of the measures conditioned on the Euclidean ball of size  $\varepsilon$  around 0 being connected to the Euclidean ball of size  $R$  around 0. The result then follows by choosing a sequence  $(\varepsilon_\delta)$  slowly tending to zero as  $\delta$  tends to zero, and then setting

$$\kappa(\delta) := \varepsilon_\delta^{-1/8} \phi_{\delta\mathbb{Z}^2, 2/3, 4}^0[0 \leftrightarrow \infty | B_{\varepsilon_\delta} \leftrightarrow \infty]$$

and

$$\Psi_{\mathcal{D}}(P; x_1, \dots, x_n) := \lim_{\varepsilon \rightarrow 0} \varepsilon^{-n/8} \mathbb{P}_{\text{CLE}(4)}(\mathcal{E}^\varepsilon(P; x_1, \dots, x_n)).$$

The asymptotics of  $\kappa(\delta)$  are due to [15, Theorem 1.2]. The conformal covariance follows easily from properties of CLE(4).  $\square$

*Proof of Corollary 1.4.* This is a direct consequence of the Edwards–Sokal coupling and Corollary 1.3.  $\square$

**AI use statement.** No mathematical content was generated or found with the assistance of artificial intelligence.

**Acknowledgments.** The second author was funded by the ERC SPaPaF. The first, second, third, and fifth authors were funded by the NCCR SwissMap and the Swiss FNS. The fourth author acknowledges support by the French National Research Agency (ANR), project number ANR-23-CPJ1-0150-01, and the project ERC GeStruct.

## References

- [1] E. Averous, H.B. Chen, H. Duminil-Copin, T. He, D. Krachun, I. Manolescu, and J. Xia, Rotational invariance of six-vertex model scaling limits for  $-1 \leq \Delta \leq -1/2$ . *preprint*, 2026.
- [2] E. Averous, H. Duminil-Copin, T. He, P. Lammers and I. Manolescu, Conformal invariance of the six-vertex model’s height function. *preprint*, 2026.
- [3] E. Averous, H. Duminil-Copin, P. Lammers, and I. Manolescu. Conformal invariance of the planar Ashkin-Teller model, *Manuscript in preparation*, 2026.
- [4] Rodney J. Baxter. *Exactly Solved Models in Statistical Mechanics*. London; New York: Academic Press, 1982.
- [5] R. J. Baxter, S. B. Kelland and F. Y. Wu, Equivalence of the Potts model or Whitney polynomial with an ice-type model. *Journal of Physics A: Mathematical and General*, 9(3) :397, 1976.
- [6] A. A. Belavin, A. M. Polyakov and A. B. Zamolodchikov, Infinite conformal symmetry in two-dimensional quantum field theory. *Nuclear Physics B*, 241(2):333–380, 1984.

- [7] A. A. Belavin, A. M. Polyakov and A. B. Zamolodchikov, Infinite conformal symmetry of critical fluctuations in two dimensions. *Journal of Statistical Physics*, 34(5-6):763–774, 1984.
- [8] S. Benoist and C. Hongler, The scaling limit of critical Ising interfaces is CLE(3). *Annals of Probability*, 47(4):2049–2086, 2019.
- [9] N. Berestycki and E. Powell, *Gaussian Free Field and Liouville Quantum Gravity*. Cambridge Studies in Advanced Mathematics. Cambridge: Cambridge University Press, 2025.
- [10] F. Camia and C.M. Newman, Two-dimensional critical percolation: the full scaling limit. *Communications in Mathematical Physics*, 268(1):1–38, 2006.
- [11] F. Camia and C.M. Newman, Critical percolation exploration path and SLE<sub>6</sub>: a proof of convergence. *Probability Theory and Related Fields*, 139(3-4):473–519, 2007.
- [12] D. Chelkak, C. Hongler and K. Izyurov, Conformal invariance of spin correlations in the planar Ising model. *Annals of Mathematics (2)*, 181(3):1087–1138, 2015.
- [13] D. Chelkak and S. Smirnov, Universality in the 2D Ising model and conformal invariance of fermionic observables. *Inventiones Mathematicae*, 189(3):515–580, 2012.
- [14] D. Chelkak, H. Duminil-Copin, C. Hongler, A. Kemppainen and S. Smirnov, Convergence of Ising interfaces to Schramm’s SLE curves. *Comptes Rendus Mathématiques*, 352(2):157–161, 2014.
- [15] H.-B. Chen, H. Duminil-Copin, T. He, F. Jacopin, D. Krachun, I. Manolescu, and J. Xia, Critical exponents for the planar random-cluster model with cluster-weight  $q = 4$ . *arXiv:2605.30030*, 2026.
- [16] H.-B. Chen, J. Xia, Multi-point functions of a full-plane two-state fuzzy 4-Potts model. *arXiv:2605.31374*, 2026.
- [17] F. David, A. Kupiainen, R. Rhodes, and V. Vargas. Liouville Quantum Gravity on the Riemann Sphere. *Communications in Mathematical Physics*, 342(3):869–907, 2016.
- [18] Julien Dubédat. Topics on Abelian Spin Models and Related Problems. *Probability Surveys*, 8:374–402, 2011.
- [19] H. Duminil-Copin, *Lectures on the Ising and Potts models on the hypercubic lattice*. In PIMS-CRM Summer School in Probability, 35–161, Springer, 2017.
- [20] H. Duminil-Copin, M. Gagnebin, M. Harel, I. Manolescu and V. Tassion, Discontinuity of the phase transition for the planar random-cluster and Potts models with  $q > 4$ . *Annales Scientifiques de l’École Normale Supérieure*, 54(6):1363–1413, 2021.
- [21] H. Duminil-Copin, M. Lis, and W. Qian. Conformal invariance of double random currents I: Identification of the limit. *Proceedings of the London Mathematical Society*, 130(1):e70022, 2025.
- [22] H. Duminil-Copin, M. Lis, and W. Qian. Conformal invariance of double random currents II: tightness and properties in the discrete. *arXiv:2107.12880*, 2021.

- [23] H. Duminil-Copin, K. Kozłowski, P. Lammers and I. Manolescu, Gaussian free field convergence of the six-vertex model with  $-1 \leq \Delta \leq -\frac{1}{2}$ . *arXiv:2603.06268*, 2026.
- [24] H. Duminil-Copin, K. Kozłowski, D. Krachun, I. Manolescu and M. Oulamara, Rotational invariance in critical planar lattice models. *arXiv:2012.11672*, 2020.
- [25] H. Duminil-Copin, J.-H. Li and I. Manolescu, Universality for the random-cluster model on isoradial graphs. *Electronic Journal of Probability*, 23:1–70, 2018.
- [26] H. Duminil-Copin, V. Sidoravicius and V. Tassion, Continuity of the phase transition for planar random-cluster and Potts models with  $1 \leq q \leq 4$ . *Communications in Mathematical Physics*, 349(1):47–107, 2017.
- [27] G. Grimmett, *The random-cluster model*, vol. 333 of Grundlehren der Mathematischen Wissenschaften, Springer-Verlag, Berlin, 2006.
- [28] C. Guillarmou, R. Rhodes, and V. Vargas. Polyakov’s Formulation of 2d Bosonic String Theory. *Publications Mathématiques de l’IHES* 130:111–185, 2019.
- [29] C. Hongler and S. Smirnov, The energy density in the planar Ising model. *Acta Mathematica*, 211(2):191–225, 2013.
- [30] R. Kenyon, Conformal invariance of domino tiling. *Annals of Probability*, 28(2):759–795, 2000.
- [31] R. Kenyon, Dominos and the Gaussian Free Field. *The Annals of Probability*, 29(3):1128–1137, 2001.
- [32] M. Khristoforov and S. Smirnov, Percolation and  $O(1)$  loop model. *arXiv:2111.15612*, 2021.
- [33] G.F. Lawler, *Conformally invariant processes in the plane*, volume 114 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2005.
- [34] G.F. Lawler, O. Schramm and W. Werner, Conformal invariance of planar loop-erased random walks and uniform spanning trees. *Annals of Probability*, 32(1B):939–995, 2004.
- [35] Elliott H. Lieb and F. Y. Wu. Two-Dimensional Ferroelectric Models. *12th School of Modern Physics on Phase Transitions and Critical Phenomena*, 1980.
- [36] J. Miller and S. Sheffield. Imaginary Geometry I: Interacting SLEs. *Probability Theory and Related Fields*, 164(3):553–705, 2016.
- [37] J. Miller and S. Sheffield. Liouville Quantum Gravity and the Brownian Map I: The QLE(8/3,0) Metric. *Inventiones mathematicae*, 219(1):75–152, 2020.
- [38] A.Z. Patashinskii and V.L. Pokrovskii, Second Order Phase Transitions in a Bose Fluid. *Soviet Physics JETP* 19(3):677–691, 1964.
- [39] A.M. Polyakov, Microscopic Description of Critical Phenomena. *Soviet Physics JETP* 28(3):533–539, 1968.

- [40] A.M. Polyakov, Properties of Long and Short Range Correlations in the Critical Region. *Soviet Physics JETP* 30(1):151–157, 1970.
- [41] A.M. Polyakov, Conformal symmetry of critical fluctuations. *Pis. ZhETP*, 12:538–541, 1970.
- [42] R.B. Potts. Some generalized order-disorder transformations. *Proceedings of the Cambridge Philosophical Society*, 48(2):106–109, 1952.
- [43] G. Ray and Y. Spinka, Finitary Codings for Gradient Models and a New Graphical Representation for the Six-Vertex Model. *Random Structures & Algorithms*, 61(1):193–232, 2022.
- [44] R. Rhodes and V. Vargas. Gaussian Multiplicative Chaos and Applications: A Review. *Probability Surveys*, 11:315–392, 2014.
- [45] Oded Schramm. Scaling Limits of Loop-Erased Random Walks and Uniform Spanning Trees. *Israel Journal of Mathematics*, 118(1):221–288, 2000.
- [46] O. Schramm and S. Sheffield, Contour lines of the two-dimensional discrete Gaussian free field. *Acta Mathematica*, 202(1):21–137, 2009.
- [47] J. Aru, A. Sepúlveda, and W. Werner. On bounded-type thin local sets of the two-dimensional Gaussian free field. *J. Inst. Math. Jussieu*, 18(3):591–618, 2019.
- [48] S. Sheffield, Gaussian free fields for mathematicians. *Probab. Theory Relat. Fields*, 139:521–541, 2007.
- [49] S. Sheffield. Exploration Trees and Conformal Loop Ensembles. *Duke Mathematical Journal*, 147(1):79–129, 2009.
- [50] S. Sheffield and W. Werner, Conformal Loop Ensembles: The Markovian Characterization and the Loop-Soup Construction. *Annals of Mathematics*, 176(3):1827–1917, 2012.
- [51] S. Smirnov, Critical percolation in the plane: conformal invariance, Cardy’s formula, scaling limits. *Comptes Rendus de l’Académie des Sciences. Série I. Mathématique*, 333(3):239–244, 2001.
- [52] S. Smirnov, Conformal invariance in random-cluster models. I. Holomorphic fermions in the Ising model. *Annals of Mathematics*, 172(2):1435–1467, 2010.